

References

Agency for Health Care Policy and Research (1995). Clinical Practice Guideline Number 16. *Post-Stroke Rehabilitation*. U.S. Department of Health and Human Services, Rockville, MD.

American College of Sports Medicine (1991) *Guideline for Graded Exercise Testing and Prescription* (4th ed.) Philadelphia: Lea & Febiger.

Anderson EM, Malgren JA (1994) Screening for depression in well older adults: evaluation of a short form of the CES-D. *American Journal of Preventive Medicine* 10(2): 77-84.

Bauman WA, Spungen MS, Zhong Y-G, Rothstein JL, Petry C, Gordon, SK (1992) Depressed serum high density lipoprotein cholesterol levels in veterans with spinal cord injury. *Paraplegia* 30; 697-703.

Bauman WA, Spungen AM, Raza M, Rothstein J, Zhang R-L, Zhong Y-G, Tsuruta M, Shahidi R, Pierson RN, Wang J, Gordon SK. Coronary artery disease: metabolic risk factors and latent disease in individuals with paraplegia. *Mount Sinai Journal of Medicine*. 1992;59:163-168.

Brenes G, Dearwater S, Shapera R, LaPorte RE, Collins E (1986). High density lipoprotein cholesterol concentrations in physically active and sedentary spinal cord injured patients. *Archives of Physical Medicine and Rehabilitation*, 67:445-450.

Broman CL (1995) Leisure-time physical activity in an African-American population. *Journal of Behavioral Medicine* 18(4):341-53.

Bruce RA, Kusumi MS, Hosmer D (1973) Maximal oxygen intake and nomographic assessment of functional aerobic impairment in cardiovascular disease. *American Heart Journal*, 85:546.

Crespo CJ, Keteyian SJ, Heath GW, Sempos CT (1996) Leisure-time physical activity among U.S. adults. Results from the Third National Health and Nutrition Examination Survey. *Archives of Internal Medicine* 156(1):93-8.

DeVivio MJ, Black, KJ, Stoner, SL (1993) Causes of death during the first twelve years after spinal cord injury. *Archives of Physical Medicine and Rehabilitation*, 74:248-254.

Drummond A (1990) Leisure after stroke. *International Disability Studies*. 12:157-160.

Duckworth WC, Solomon SS, Jallepalli P, Heckmeyer C, Finnern J, Powers A (1980) Glucose intolerance due to insulin resistance in patients with spinal cord injuries. *Diabetes* 29:906-910.

Ellestad MH (1994) *Stress Testing. Principles and Practice*, Philadelphia: F.A. Davis Company, 4th edition.

Fletcher BJ, Dunbar SB, Felner JM, et al. (1994) Exercise testing and training in physically disabled men with clinical evidence of coronary artery disease. *American Journal of Cardiology*, 73:170-4.

Folson AR, Jacobs DR, Caspersen CJ, Gomez-Marín O, Knudsen J (1986) Test-retest reliability of the Minnesota leisure time physical activity questionnaire. *Journal of Chronic Disease* 39:505-511.

Fulg-Meyer, AR, Branholm, IB, Fulg-Meyer, KS (1991) Happiness and domain-specific life satisfaction in adult northern Swedes. *Clinical Rehabilitation*, 5:25-33.

Gresham GE, Phillips TF, Wolf PA, McNamara PM, Kannel WB, Dawber TR, (1979) Epidemiologic profile of long-term stroke disability; the Framingham study. *Archives of Physical Medicine and Rehabilitation*, 60:487-91.

Hjeltnes N, Vokac Z (1979) Circulatory strain in everyday life of paraplegics *Scandinavian Journal of Rehabilitation Medicine*. 11:67-73.

Jongbloed L, Morgan M (1991) An investigation of involvement in leisure activities after a stroke. *American Journal of Occupational Therapy*. 45:420-427.

Kennedy DW, Smith RW (1990) A comparison of past and future leisure activity participation between spinal cord injured and non-disabled persons. *Paraplegia* 28:130-136.

Leblanc J, Nadeau A, Richard D (1981) Studies on the sparing effect of exercise on insulin requirements in human subjects. *Metabolism* 30:1119-24.

Paffenberger RS Jr, Hyde RT, Wing AL, Lee IM, Jung DL, Kampert JB. The association of changes in physical activity level and other lifestyle characteristics with mortality among men. *New England Journal of Medicine* 1993;328:538-545.

Peiris AN, Sothmann MS, Hoffman RG, et al. (1989) Adiposity, fat distribution, and cardiovascular risk. *Annals of Internal Medicine*, 110:867-72.

Potempa, K, Lopez, M et. al. Physiological Outcomes of Aerobic Exercise Training in Hemiparetic Stroke Patients. *Stroke*. 1995;26(1):101-105.

Reeves RA (1995) Does this patient have hypertension? How to measure blood pressure. *Journal of the American Medical Association*, 273:1211-18.

Roth, EJ, Mueller, K, Green, D (1988) Stroke rehabilitation outcome: impact of coronary artery disease. *Stroke* Jan 19(1):42-7.

Roth EJ, (1993) Heart disease in patients with stroke: incidence, impact, and implications for rehabilitation part I: classification and prevalence. *Archives of Physical Medicine and Rehabilitation*, 74:752-760.

Stampfer MJ , Sacks M , Salvini S, Willett WC , Hennekens, C H. A prospective study of cholesterol, apolipoproteins, and the risk of myocardial infarction *New England Journal of Medicine*. 1991;325:373-381.

Stover SL , DeLisa JA , Whitneck GG (Eds.) (1995) *Spinal Cord Injury: Clinical Outcomes from the Model Systems*. Aspen Publishers, Inc., Gaithersburg, Maryland

Sullivan LW, (1991) Striking out stroke in America. *Be Stroke Smart* 9(2):1.

Taylor HL, Jacobs DR, Schucker B, Knudsen J, Leon AS, Debacker G (1978) A questionnaire for the assessment of leisure time physical activities. *Journal of Chronic Disease* 31:741-755.

Ware J & Sherbourne C. The MOS 36 item Short Form Health Survey (SF-36) *Medical Care*. 1992;30:473-483.